

①

NPS-PH-91-004

# NAVAL POSTGRADUATE SCHOOL

Monterey, California

AD-A232 222



DTIC  
ELECTE  
FEB 26 1991  
S B D

ATMOSPHERIC OPTICAL TURBULENCE MEASUREMENTS  
TAKEN AT  
ANDERSON MESA, FLAGSTAFF, ARIZONA  
BETWEEN 13-19 NOVEMBER 1989

by

G. Tirrell Vaucher, C.A. Vaucher,  
and D.L. Walters

Approved for public release; distribution unlimited

Prepared for: Naval Research Laboratory  
Washington, D.C. 20375

WL/ARCA  
Kirtland Air Force Base  
New Mexico 97117

91 2 22 046

NAVAL POSTGRADUATE SCHOOL  
Monterey, California


Rear Admiral R.W. West, Jr.  
Superintendent


H. Shull  
Provost


The work reported herein was supported in part by the Naval Research Laboratory Program at the Naval Postgraduate School with funds provided by the Naval Research Laboratory, Air Force Weapons Laboratory, and the Naval Postgraduate School.

Reproduction of all or part of this report is authorized.

This report was prepared by:


  
GAIL TIRRELL VAUCHER  
Physical Scientist

  
CHRISTOPHER A. VAUCHER  
Visiting Scientist,  
Universities Space  
Research Association

  
DONALD L. WALTERS  
Associate Professor  
of Physics

Approved by:

Released by:

  
KARLHEINZ E. WOEHLE  
Chairman of Physics

  
PAUL J. MARTO  
Dean of Research

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No 2704-0188	
1a REPORT SECURITY CLASSIFICATION Unclassified			1b RESTRICTIVE MARKING		
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		
2b DECLASSIFICATION/DOWNGRADING SCHEDULE					
4 PERFORMING ORGANIZATION REPORT NUMBER NPS-PH-91-004			5 MONITORING ORGANIZATION REPORT NUMBER NPS-PH-91-004		
6a NAME OF PERFORMING ORGANIZATION Naval Postgraduate School		6b OFFICE SYMBOL (if applicable) PH		7a NAME OF MONITORING ORGANIZATION Naval Research Laboratory	
6c ADDRESS (City, State, and ZIP Code) Monterey, California 93943-5000			7b ADDRESS (City, State, and ZIP Code) Washington, D.C. 20375		
8a NAME OF FUNDING SPONSORING ORGANIZATION Naval Research Lab WL/ARCA(Air Force)		8b OFFICE SYMBOL (if applicable)		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER MIPR# WLK-0-143	
8c ADDRESS (City, State, and ZIP Code) Naval Research Lab. WL/ARCA (Air Force) Washington, D.C. Kirtland AFB 20375 New Mexico 87117			10 SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO PROJECT NO TASK NO WORK UNIT ACCESSION NO		
11 TITLE (Include Security Classification) ATMOSPHERIC OPTICAL TURBULENCE MEASUREMENTS TAKEN AT ANDERSON MESA, FLAGSTAFF, ARIZONA BETWEEN 13-19 NOVEMBER 1989 (Unclassified)					
12 PERSONAL AUTHOR(S) Gail Tirrell Vaucher, Christopher A. Vaucher, Donald L. Walters					
13a TYPE OF REPORT Technical Report		13b TIME COVERED FROM TO		14 DATE OF REPORT (Year, Month, Day) 1991 January 11	
				15 PAGE COUNT 76	
16 SUPPLEMENTARY NOTATION The view expressed in this technical report are those of the authors and do not reflect the official policy or position of the Dept of Defense or the US government.					
17 COSATI CODES FIELD GROUP SUB-GROUP			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number) atmospheric optical turbulence, transverse coherence length, isoplanatic angle, $\star H$		
19 ABSTRACT (Continue on reverse if necessary and identify by block number)  From 13 to 19 November 1989, the Naval Postgraduate School Atmospheric Optics Group acquired atmospheric optical turbulence measurements at the 31-inch Lowell Observatory telescope dome facility on Anderson Mesa, 16 km southeast of Flagstaff, Arizona. The parameters measured, the transverse coherence length and the isoplanatic angle, were part of an ongoing site survey for a large-scale, ground-based, synthetic aperture system (100-300 m baseline stellar interferometer). This report compiles, analyses and summarizes the acquired optical data. Also discussed are the synoptic meteorological events present during the data acquisition period.					
20 DISTRIBUTION AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> OTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a NAME OF PERSONAL AUTHOR(S) Gail Tirrell Vaucher			22b TELEPHONE (Include Area Code) (22c DDD OFFICE USE) (408)-646-3207/2267 PH (PH/We)		

ATMOSPHERIC OPTICAL TURBULENCE MEASUREMENTS

TAKEN AT

ANDERSON MESA, FLAGSTAFF, ARIZONA

BETWEEN 13-19 NOVEMBER 1989

by

G.Tirrell Vaucher, C.A. Vaucher, and D.L. Walters

Atmospheric Optics Group  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93943-5000

# ABSTRACT

From 13 to 19 November 1989, the Naval Postgraduate School Atmospheric Optics Group acquired atmospheric optical turbulence measurements at the 31-inch Lowell Observatory telescope dome facility on Anderson Mesa, 16 km southeast of Flagstaff, Arizona. The parameters measured, the transverse coherence length and isoplanatic angle, were part of an ongoing site survey for a large-scale, ground-based, synthetic aperture system (100-300 m baseline stellar interferometer). This report compiles, analyses and summarizes the acquired optical data. Also discussed are the synoptic meteorological events present during the data acquisition period.



<b>Accession For</b>	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
<b>Availability Codes</b>	
Dist	Avail and/or Special
A-1	

## TABLE OF CONTENTS

I.	INTRODUCTION .....	1
II.	EXPERIMENT OVERVIEW .....	3
	A. SITE TOPOGRAPHY .....	3
	B. DATA ACQUISITION .....	3
	1. Optical Instrumentation and Measurements....	3
	2. Meteorological Surface Data .....	5
	3. Synoptic Weather Information .....	5
III.	DATA ANALYSIS .....	6
	A. OPTICAL DATA ANALYSIS .....	6
	1. Transverse Coherence Length Data, $r_0$ .....	6
	2. Isoplanatic Angle Data, $\theta_0$ .....	10
	B. GENERAL SYNOPTIC WEATHER REVIEW .....	14
IV.	DATA SUMMARY .....	15
	A. CONSOLIDATING THE PARAMETERS .....	15
	B. OPTICAL/METEOROLOGICAL DATA CORRELATIONS .....	16
V.	CONCLUSIONS/RECOMMENDATIONS .....	19
APPENDIX A.	DAILY SYNOPTIC WEATHER SUMMARY .....	20
APPENDIX B.	NWS 850 AND 200 MB SYNOPTIC WEATHER CHARTS	23
APPENDIX C.	PROCESSED OPTICAL DATA (1989 NOVEMBER 13-19) .....	37
APPENDIX D.	TRANSVERSE COHERENCE LENGTH STATISTICS ....	45

APPENDIX E. ISOPLANATIC ANGLE STATISTICS .....	53
APPENDIX F. SEPTEMBER/NOVEMBER $r_0$ AND $\theta_0$ DISTRIBUTION..	61
LIST OF REFERENCES .....	64
INITIAL DISTRIBUTION LIST .....	65

## List of Tables

Table 1.	TRANSVERSE COHERENCE LENGTH STATISTICS .....	7
Table 2.	ISOPLANATIC ANGLE STATISTICS .....	11
Table 3.	OPTICAL/METEOROLOGICAL SUMMARY - ANDERSON MESA, AZ .....	15



## List of Figures

Fig 1.	Topographical Views of the Lowell Sites, Anderson Mesa, Arizona .....	4
Fig 2.	Average Transverse Coherence Lengths (89 Nov 13-19) .....	7
Fig 3.	Normalized Transverse Coherence Length Frequency Distributions .....	8
Fig 4.	Cumulative Normalized $r_0$ Frequency Distribution	9
Fig 5.	Average Isoplanatic Angles (89 Nov 13-19) .....	11
Fig 6.	Normalized Isoplanatic Angle Frequency Distributions .....	12
Fig 7.	Cumulative Normalized $\theta_0$ Frequency Distribution	13
Fig 8.	NWS 850 and 200 mb Charts: November 13, 0000 UTC	24
Fig 9.	NWS 850 and 200 mb Charts: November 13, 1200 UTC	25
Fig 10.	NWS 850 and 200 mb Charts: November 14, 0000 UTC	26
Fig 11.	NWS 850 and 200 mb Charts: November 14, 1200 UTC	27
Fig 12.	NWS 850 and 200 mb Charts: November 15, 0000 UTC	28
Fig 13.	NWS 850 and 200 mb Charts: November 15, 1200 UTC	29
Fig 14.	NWS 350 and 200 mb Charts: November 16, 0000 UTC	30
Fig 15.	NWS 850 and 200 mb Charts: November 17, 0000 UTC	31
Fig 16.	NWS 850 and 200 mb Charts: November 17, 1200 UTC	32
Fig 17.	NWS 850 and 200 mb Charts: November 18, 0000 UTC	33
Fig 18.	NWS 850 and 200 mb Charts: November 18, 1200 UTC	34
Fig 19.	NWS 850 and 200 mb Charts: November 19, 0000 UTC	35
Fig 20.	NWS 850 and 200 mb Charts: November 19, 1200 UTC	36
Fig 21.	Anderson Mesa, Az Optical Data: 1989 November 13	38
Fig 22.	Anderson Mesa, Az Optical Data: 1989 November 14	39
Fig 23.	Anderson Mesa, Az Optical Data: 1989 November 15	40
Fig 24.	Anderson Mesa, Az Optical Data: 1989 November 16	41
Fig 25.	Anderson Mesa, Az Optical Data: 1989 November 17	42
Fig 26.	Anderson Mesa, Az Optical Data: 1989 November 18	43
Fig 27.	Anderson Mesa, Az Optical Data: 1989 November 19	44
Fig 28.	Anderson Mesa, Az $r_0$ Statistics: 1989 Nov 13 ...	46
Fig 29.	Anderson Mesa, Az $r_0$ Statistics: 1989 Nov 14 ...	47
Fig 30.	Anderson Mesa, Az $r_0$ Statistics: 1989 Nov 15 ...	48
Fig 31.	Anderson Mesa, Az $r_0$ Statistics: 1989 Nov 16 ...	49
Fig 32.	Anderson Mesa, Az $r_0$ Statistics: 1989 Nov 17 ...	50
Fig 33.	Anderson Mesa, Az $r_0$ Statistics: 1989 Nov 18 ...	51
Fig 34.	Anderson Mesa, Az $r_0$ Statistics: 1989 Nov 19 ...	52

Fig 35.	Anderson Mesa, Az $\theta_0$ Statistics: 1989 Nov 13 ...	54
Fig 36.	Anderson Mesa, Az $\theta_0$ Statistics: 1989 Nov 14 ...	55
Fig 37.	Anderson Mesa, Az $\theta_0$ Statistics: 1989 Nov 15 ...	56
Fig 38.	Anderson Mesa, Az $\theta_0$ Statistics: 1989 Nov 16 ...	57
Fig 39.	Anderson Mesa, Az $\theta_0$ Statistics: 1989 Nov 17 ...	58
Fig 40.	Anderson Mesa, Az $\theta_0$ Statistics: 1989 Nov 18 ...	59
Fig 41.	Anderson Mesa, Az $\theta_0$ Statistics: 1989 Nov 19 ...	60
Fig 42.	Cumulative $r_0$ Distribution: 1989 Sept and Nov ..	62
Fig 43.	Cumulative $\theta_0$ Distribution: 1989 Sept and Nov ..	63

#### ACKNOWLEDGEMENTS

We would like to express our sincere appreciation to the Lowell Observatory, particularly Drs. Nat White and Bob Millis (Director), for their assistance during the measurement session and for the use of their 31" telescope dome facility on Anderson Mesa. Also, a special thank you to Nancy Alexander, from Technical and Business Systems, for her many long hours of support with the data acquisition.

## I. INTRODUCTION

From 13 to 19 November 1989, the Naval Postgraduate School (NPS) Atmospheric Optics Group acquired atmospheric optical turbulence measurements at the 31-inch Lowell Observatory telescope dome on Anderson Mesa, 16 km southeast of Flagstaff, Arizona. This ensemble of transverse coherence lengths ( $r_0$ ) and isoplanatic angles ( $\theta_0$ ) was part of an ongoing site survey mission for the Naval Research Laboratory large-scale, ground-based, synthetic aperture system (100-300 m baseline stellar interferometer) and for the NPS-US Air Force optimal site characterization program.

The purpose of this report is three-fold: to document the November 1989 Anderson Mesa optical measurements; to provide a statistical analysis and summary of these optical data; and, to correlate the meteorological activities (using synoptic weather charts) with the optical conditions present during the experiment period. Due to the considerable scale-differential between synoptic weather phenomena (kilometers) and optical turbulence producing events (meters), only a preliminary investigation into the third objective was undertaken. Any detailed correlations would require meso- and micro-scale meteorological information (e.g., high resolution rawinsonde measurements). Such resources were unavailable for this measurement session.

Six appendices supplement this report:

Appendix A presents daily summaries of the synoptic weather conditions prevalent over Anderson Mesa during the optical data acquisition period.

Appendix B contains hand-edited reproductions of the National Weather Service (NWS) 850 and 200 mb synoptic weather charts for the period 13-19 November 1989. For reference, important meteorological features from the NWS surface maps are superimposed onto the 850 mb charts.

Appendix C displays all processed  $r_0$  and  $\theta_0$  data sampled between 13-19 November 1989 Universal Time Coordinated (UTC). Each figure displays a separate night-time session.

Appendices D and E provide nightly, un-normalized percent frequency distributions and empirical seeing quality histograms for  $r_0$  and  $\theta_0$ , respectively. Specific bin intervals for the  $r_0$  and  $\theta_0$  distributions, as well as the empirically derived seeing quality intervals, are listed at the start of each Appendix.

Appendix F presents a cumulative September/November 1989 normalized frequency distribution for both  $r_0$  and  $\theta_0$ . The measurements incorporated in these figures represent all the 17-28 September and 13-19 November 1989 processed NPS optical data taken at both Anderson Mesa and the United States Naval Observatory, Flagstaff, Arizona.

## II. EXPERIMENT OVERVIEW

### A. SITE TOPOGRAPHY

As described in Vaucher, Vaucher and Walters (1990), Anderson Mesa is an 125 m high plateau situated in the ponderosa pine and lake mesa-country 16 km southeast of Flagstaff and 18 km west of the high desert floor. The 31-inch telescope dome used for optical data gathering is 2.2 km above sea level and located on the southwest edge of the mesa. Figure 1 displays a 3-dimensional topographical view of Anderson Mesa, showing the location of the 31-inch site, as well as major features of interest. The contour interval is 5 meters.

### B. DATA ACQUISITION

All data acquisition sessions commenced at local sunset and concluded with the onset of local sunrise twilight. The total sampling duration was approximately 10-11 hours per night. Intermittent cloud-cover often disrupted the ability to sample data, providing both gaps in the individual nightly optical records, as well as severely restricting the total volume of measurements taken during some sessions.



#### 1. Optical Instrumentation and Measurements

The optical turbulence parameters gathered throughout the experiment include the isoplanatic angle ( $\theta_0$ ) and the transverse coherence length ( $r_0$ ). A brief description of these parameters is available in Vaucher (1989). The isoplanometer and transverse coherence length sensors were designed and built by Dr. D.L. Walters. Stevens (1985) and Walters, Favier, and Hines (1979), respectively describe specific details for each instrument. Vaucher, Vaucher and Walters (1990) explain the optical system, configurations, and data acquisition procedures utilized during the Anderson Mesa site survey missions.

All optical measurements were recorded in Universal Time Coordinated (UTC). The conversion from local Mountain Standard Time (MST) to UTC is:

$$\text{Time(UTC)} = \text{Time(MST)} + 7 \text{ hours.}$$

Arizona retains MST throughout the year.



**Fig 1. Topographical Views of the Lowell Sites, Anderson Mesa, Arizona**

## 2. Meteorological Surface Data

A model 5165-A WeatherMeasure (WEATHERtronics) Humidity/Temperature Indicator periodically measured the surface meteorological thermodynamic data. Vaucher, Vaucher and Walters (1990) present the detailed specifications for this device. The main function of this probe was to monitor the multiple weather changes and potentially hazardous (extremely cold) environments during the data acquisition.

## 3. Synoptic Weather Information

GOES-WEST Visible Satellite Images, as well as surface, 500 mb, and 200 mb National Weather Service charts provided an "on-site" evaluation of the synoptic weather conditions. These also helped to detect trends and potential sources of optical turbulence during the experiment period.

Post-experiment analyses of synoptic weather activity around Flagstaff were based on the six National Weather Service standard isobaric charts. These constant pressure surfaces and their equivalent heights are:

<u>Pressure (mb)</u>	<u>Equivalent Height Above Sea Level (km)</u>
Surface	0.1
850	1.4
700	3.0
500	5.5
300	9.2
200	12.0

It should be noted that the equivalent heights indicated above represent average values. For any given pressure level, the actual height will vary as Low and High pressure systems traverse the site (Vaucher, Vaucher, and Walters, 1990). Appendix B provides two of the six NWS isobaric surfaces (850 and 200 mb) used in post-analysis.



### III. DATA ANALYSIS

#### A. OPTICAL DATA ANALYSIS

##### 1. Transverse Coherence Length Data, $r_o$

Using the Appendix D  $r_o$  Empirical Seeing Quality Scale, the dominant  $r_o$  optical condition for 13-19 November is mediocre (51-100 mm). Figure 2 displays the oscillatory pattern of average (and standard deviation)  $r_o$  as a function of the observing date. Values plotted in this figure are listed in Table 1, along with the nightly  $r_o$  standard deviation of the mean. Minimum and maximum average  $r_o$  data occur on 15 and 17 November, respectively. November 17 also exhibits the greatest standard deviation of the mean (2.5 mm), but this can be explained: examining the processed data for this date (Appendix C), the initial 5-6 hours of  $r_o$  samples are between 51-100 mm; the remaining 5-6 hours are significantly higher (75-275 mm). In contrast, the large standard deviation of the mean calculated for 16 November (2.3 mm) is a function of the limited number of points collected (65 samples).

Figure 3 presents the normalized  $r_o$  frequency distributions for each session. These curves generally peak at 100 mm or less, though values above 100 mm are present in all but the 18 November data. The session with the greatest frequency of "good" to "very good"  $r_o$  (100 mm and greater) is 17 November 1989 (Table 3).

Compiling all  $r_o$  samples acquired between 13-19 November, the cumulative normalized  $r_o$  frequency distribution (Figure 4) indicates that 70-80 mm is the interval with the greatest number of samples. Nearly two-thirds (64 %) of all measurements taken have values between 51 and 100 mm ("mediocre").

TABLE 1. TRANSVERSE COHERENCE LENGTH STATISTICS

Date (UTC)	Number of Data Points	Average $r_0$ (mm)	Standard Deviation (mm)	Standard Deviation of Mean (mm)
13 Nov	430	91.6	22.7	1.1
14 Nov	410	77.5	18.3	0.9
15 Nov	351	48.7	13.9	0.7
16 Nov	65	89.3	18.4	2.3
17 Nov	368	101.2	47.3	2.5
18 Nov	241	51.3	11.5	0.7
19 Nov	238	71.2	14.0	0.9





Fig 2. Average Transverse Coherence Lengths (89 Nov 13-19)  
Solid line is data average; Dashed line is standard  
deviation of the data.

NORMALIZED  $r_0$  FREQUENCY DISTRIBUTION  
Anderson Mesa, AZ - 1989 November 13-16



NORMALIZED  $r_0$  FREQUENCY DISTRIBUTION  
Anderson Mesa, AZ - 1989 November 17-19




Fig 3. Normalized Transverse Coherence Length Frequency Distributions for Anderson Mesa, AZ (1989 Nov 13-19).

CUMULATIVE NORMALIZED  $r_0$  FREQUENCY DISTRIBUTION  
Anderson Mesa, AZ - 1989 November 13-19




Fig 4. Cumulative Normalized  $r_0$  Frequency Distribution  
for the Anderson Mesa 1989 Nov 13-19 session.

## 2. Isoplanatic Angle Data, $\theta_0$

Figure 5 shows that the average nightly isoplanatic angles ( $\theta_0$ ) also display an oscillatory pattern. The first order trend for the seven night period is toward smaller angles (more turbulent atmosphere). The average  $\theta_0$  values (Table 2) begin around 12 urad (13-14 November), then decrease to nearly 6 urad (19 November).  $\theta_0$  minima occur on 15 and 18 November, with three maxima occurring on 14, 16 and 19 November. While most values are "poor" to "mediocre" (4-12 urad), five of the seven nights display "good" (12-20 urad) optical numbers (see Appendix E for the  $\theta_0$  Empirical Seeing Quality Scale). The greatest frequency of "good" values (49%) appears on 13 November 1989 (Table 3).

Figure 6 presents the normalized  $\theta_0$  frequency distributions for each session. The maxima for these curves occur between 3-13 urad. Individual values above 12 urad ("good" to "excellent") are present on 13, 14, and 17 November. The session with the greatest frequency of "good" to "excellent"  $\theta_0$  is 13 November (Table 3).

Compiling all 13-19 November samples, the cumulative normalized  $\theta_0$  frequency distribution (Figure 7) has a primary maximum of 4-5 urad, with secondary peaks around 11 and 13 urad. Two-thirds (66%) of all cumulative  $\theta_0$  samples fall within the range of 0-8 urad ("very poor" to "poor").

TABLE 2. ISOPLANATIC ANGLE STATISTICS

Date (UTC)	Number of Data Points	Average $\theta_0$ (urad)	Standard Deviation (urad)	Standard Deviation of Mean (urad)
13 Nov	1029	11.86	1.80	0.06
14 Nov	3643	11.94	2.58	0.04
15 Nov	3129	4.45	0.95	0.02
16 Nov	63	8.46	0.73	0.09
17 Nov	2914	6.07	4.34	0.08
18 Nov	1780	3.19	0.99	0.02
19 Nov	1636	6.19	1.27	0.03





Fig 5. Average Isoplanatic Angles (89 Nov 13-19) - Solid line is data average; Dashed line is standard deviation of the data.

NORMALIZED  $\theta$ . FREQUENCY DISTRIBUTION  
Anderson Mesa, AZ - 1989 November 13-16



NORMALIZED  $\theta$ . FREQUENCY DISTRIBUTION  
Anderson Mesa, AZ - 1989 November 17-19




Fig 6. Normalized Isoplanatic Angle Frequency Distributions for Anderson Mesa, AZ (89 Nov 13-19).




Fig 7. Cumulative Normalized  $\theta_0$  Frequency Distribution for the Anderson Mesa 1989 Nov 13-19 session.



## B. GENERAL SYNOPTIC WEATHER REVIEW

The overall synoptic weather pattern for the 13-19 November can be categorized into events occurring within three layers:

From the surface to 700 mb, the northern Pacific High expands eastward over Nevada/Utah. Meanwhile, shortwaves (850- 700 mb) propagate through a mid-continental trough just east of the northern Pacific High.

On the 500 mb level, the site is influenced by the northwesterly circulation associated with the western edge of a mid-continental trough.

Aloft (300-200 mb), polar and subtropical jets (optical turbulence sources) near and overrun the site.

For a day by day review of the synoptic weather activity, see Section IV (Data Summary) and Appendix A. Appendix B provides NWS charts for additional reference.

# IV. DATA SUMMARY

## A. CONSOLIDATING THE PARAMETERS

Table 3 consolidates key optical and meteorological parameters acquired between 13 and 19 November 1989. While the optical mean values (Parts A and B) appear earlier, the "Empirical % >='Good'" columns are seen for the first time. Here, the Empirical Seeing Quality Scale (Appendices D and E) quantifies the percentage of the individual datum points within an observing session that fall into the "good" or better classifications. The "Empirical Dominant Conditions" columns provide a quick interpretation of typical  $r_0$  and  $\theta_0$  numbers for each night, as calibrated by the empirical seeing quality scale.

**TABLE 3. OPTICAL/METEOROLOGICAL SUMMARY-ANDERSON MESA, AZ**

A. Transverse Coherence Length:				B. Isoplanatic Angle:			
Nov	Mean	Empirical:		Mean	Empirical:		
Date	Value	Dominant	% >=	Value	Dominant	% >=	
(UTC)	(mm)	Conditions	"Good"	(urad)	Conditions	"Good"	
13	91.6	Mediocre(74%)	25	11.86	Medio/Good(49%)	49	
14	77.5	Mediocre(89%)	8	11.94	Good(48%)	48	
15	48.7	Poor(69%)	1	4.45	Poor(70%)	0	
16	89.3	Mediocre(74%)	26	8.46	Mediocre(68%)	0	
17	101.2	Mediocre(46%)	46	6.07	Poor(65%)	8	
18	51.3	Poor(51%)	0	3.19	Very Poor(78%)	0	
19	71.2	Mediocre(93%)	3	6.19	Poor(91%)	0	

## C. Synoptic Meteorological Conditions:

Nov	Maximum Wind Speed Gradient		Frontal(F)/
Date	Horizontal	Wind Shear (at 200 mb)	Non-Frontal(NF)/
(UTC)		( $\times 10^{-3}$ ) $s^{-1}$	Transitional(T)
13		2.6	T
14		1.8	T
15		1.2	F
16		1.7	T
17		2.9	T
18		6.4	T
19		2.1	T

The presence of 200 and 300 mb jet stream flow (70 kt and greater) over a site has been loosely employed as an indicator of turbulence aloft (where  $\theta_0$  is especially sensitive). To quantify the jet stream activity, the 200 mb NWS synoptic weather chart isotaches were used to calculate the maximum wind speed gradient (200 mb Horizontal Wind Shear) over the site. These results are tabulated in Part C of Table 3.

Another column in the meteorological parameters section (Part C), the "Frontal/Non-Frontal/Transitional", aids in interpreting the potential optical quality over the night. Frontal and Non-frontal weather are as defined in Section IV of Vaucher, Vaucher, and Walters (1990). Generally, Frontal weather produces poor optical conditions while Non-Frontal weather renders a stable atmosphere of generally high optical quality. Transitional implies a changeover period from either Frontal or Non-Frontal conditions. It often signifies an atmosphere in which separate cold and warm air masses collide and mix, frequently producing very turbulent layers and subsequently poor optical seeing.

## B. OPTICAL/METEOROLOGICAL DATA CORRELATIONS

The following discussion assimilates the various important elements presented in the Data Analysis Section (Part III) and, Appendices A through E. Table 3 should be referenced throughout.

The initial two nights (13-14 November) of optical data collection were predominantly "mediocre" for both  $r_0$  and  $\theta_0$ . Synoptic weather activity over this time period included a turbulence-inducing shortwave in the 850-700 mb pressure layer, as well as a mid-to-upper level (500-200 mb pressure layer) Low centered over the Mexico/Arizona border. Wind shear aloft was magnified with the approach of accelerated airflows (70 knot jets) to the north and south.

The smallest  $r_0$  and  $\theta_0$  values measured (most turbulent conditions) were sampled on 15 November 1989. Initially (0100 UTC),  $r_0$  were centered around 90 mm. Later that night, the values decreased to approximately 40 mm (0400 to 1230 UTC). The  $\theta_0$  values went from poor (4-8 urad) to very poor (0-4 urad) during the same time period. The overall wretched optical conditions were a product of a variety of turbulence producing events: a cold front traversing the

site; a transition from an eastward moving Low to the clockwise circulation of the northern Pacific High spreading over Nevada; and, an approaching 70-110 kt jet aloft.

Seeing conditions, as measured by  $r_o$ , gradually improved over the next two nights (16-17 November). Note that on 16 November, sampling was severely restricted due to an extensive cloud cover. Hence, statistics derived from this limited session should be cautiously considered.

On 17 November, 46% of the  $r_o$  samples measured were between 101-300 mm ("good" to "very good"), 46% were between 51-100 mm ("mediocre") and 8% were between 0-50 mm ("poor") (Appendix D). The  $\theta_o$  values indicated turbulence aloft: 8% were between 12-20 urad ("good"), while the remaining 92% were between 0-11 urad ("very poor" to "mediocre") (Appendix E). The transitional synoptic events producing these contrasting  $r_o$  and  $\theta_o$  measurements included: an eastern extension of the northern Pacific High competing with a Low over Mexico. During this period, the High dominated. Aloft, however, a 70-110 kt jet flowed counterclockwise within 5-8 degrees latitude (555-888 km) of the site. Though synoptic weather charts indicate that the jet did not traverse the site, the shear set up by the proximity of the jet flow was enough to suppress the isoplanatic angles.

In the last two sessions (18-19 November),  $r_o$  dipped to an average value of 51 and 71 mm (both "mediocre"), respectively. The corresponding average  $\theta_o$  were 3 and 6 urad ("very poor"/"poor"). The synoptic weather scenario for 18 November continued to be transitional (Table 3). The site was under a southeastward moving surface High centered in the northwestern states and a surface Low over Mexico. At 850 mb, the site was on the edge of the northern Pacific High extension and a flow through the mid-continental trough. From 700-500 mb, an eastward moving shortwave placed the site under a weak Low. This Low became more organized between 0000 and 1200 UTC. Aloft, a 70 to 110 kt jet initially (0000 UTC) circumnavigated the site, passing within 5-8 degrees latitude in all directions. By 1200 UTC, a 115 kt northwesterly jet passed over the mesa at 200 mb. Wind shear for the 200 mb level began at  $4.3E-5 \text{ s}^{-2}$  (0000 UTC) and increased to  $8.5E-5 \text{ s}^{-2}$  (very strong shear!!) by 1200 UTC. This pattern of strong wind shear often results in low optical turbulence numbers. The 18 November measurements reinforced this general observation.

On 19 November 1989, the synoptic weather pattern over the site included a High pressure system associated with the northern Pacific High extending over Utah. The presence of this High was observed from the surface to 500 mb. At 500 mb, the site bordered the High just described above, and a Low over Baja, California. Aloft, the site was directly under a jet supporting a Low over Mexico. During the 0000-1200 UTC time interval, this upper level Low appeared to develop as a cut-off Low over Mexico/southwestern states. Wind shear over the site at 200 mb was about  $2.1\text{E-}5 \text{ s}^{-1}$ ; slightly weaker than 18 November, yet still a significant element for optically turbulent conditions aloft (low  $\theta_0$  values).

The greatest 200 mb horizontal wind shear occurred on 18 November. Also on this date, the mean  $\theta_0$  was the smallest magnitude of the seven day session, with dominant conditions empirically classed as "very poor" (78%). The 18 November mean  $\tau_0$  likewise registered a low magnitude (51.3 mm), with the dominant conditions rated as "poor" (51%). Unfortunately, the least turbulent conditions measured by the optical parameters do not coincide with the smallest 200 mb wind speed gradient. Since optical turbulence is a function of both the three-dimensional wind shear and temperature gradient integrated through an entire atmospheric cross-section, a good correlation between a single level horizontal wind shear (e.g., 200 mb) and the general optical conditions cannot be expected.

## V. CONCLUSIONS/RECOMMENDATIONS

The primary purpose of this study was to evaluate the Flagstaff, Arizona region as a potential site for a large baseline stellar interferometer. The initial measurement session completed in September 1989 by the Atmospheric Optics Group (NPS) found that "despite the constant colliding of cold and warm air masses typical for this area in September, significant Good to Excellent seeing conditions occurred". (Vaucher, Vaucher, Walters, 1990)

In contrast to September 1989, the November 1989 measurement session yielded only limited "good" quality optical data. The fragmented layers of contrasting air masses (a predominantly transitional weather pattern), coupled with an accelerated circulation (jet) over the site (300-200 mb), rendered an almost continuously turbulent optical environment. A glimmer of favorable atmospheric optical conditions occur on 13-14 November for  $\theta_0$  and 17 November for  $r_0$ . Because the data discussed throughout the previous sections are only a snapshot of the true atmospheric optical conditions over Anderson Mesa, the original September 1989 site report recommendation remains: for an ideal optical database, one should take samples every night for several years. More realistically, the authors propose acquiring a seasonal 7-10 night optical dataset coupled with a climatological study of the site. To attain a better understanding of the events producing the measured optical conditions, a minimum of three on-site, high resolution, rawinsonde launches per observation night are recommended.

In summary, the Flagstaff region warrants further consideration. It must be examined seasonally, for at least one to two years more before crude conclusions as to its optical viability can be determined.

## APPENDIX A. DAILY SYNOPTIC WEATHER SUMMARY

Site: Anderson Mesa, Flagstaff, Arizona  
Time Period: 13-19 November 1989 (UTC)  
Equipment Used: Transverse Coherence Length Sensor  
Isoplanatic Angle Sensor  
National Weather Service Synoptic Charts

The overall synoptic weather pattern for the 13-19 November 1989 time period consists of:

surface-700 mb: The northern Pacific High expanding eastward over Nevada/Utah;  
850-700 mb: Shortwaves propagating through a mid-continental trough;  
500 mb: The western edge of a mid-continental, longwave trough dictating the circulation pattern;  
300-200 mb: Polar/subtropical jets (source of optical turbulence) nearing/overrunning the site.

The "worst" seeing was sampled on 15 November 1989. Transverse coherence lengths began around 90 mm (0100 UTC), then decreased to approximately 40 mm (0400 to 1230 UTC). Isoplanatic angles went from poor (4-8 urad) to very poor (0-4 urad) during the same time period. The overall wretched optical conditions were a product of a cold front traversing the site coupled with an approaching 70-110 knot (kt) jet aloft.

The "best" seeing was measured on 17 November 1989. Transverse coherence lengths were generally between 100-250 mm from 0600 to 1045 UTC. The isoplanatic angles during this time period, however, were poor (4-8 urad). The major synoptic events creating these apparently dichotomous optical measurements may be explained by the presence of a High pressure center dominating conditions, with a (wind) jet aloft. Specifically, an eastern extension of the northern Pacific High was competing with a Low over Mexico. During this session, the High dominated. Aloft, however, a 70-110 kt jet flowed counterclockwise within 5-8 degrees latitude (555-888 km) of the site. Though the synoptic weather charts indicate that the jet does not traverse the site, the shear set up by the proximity of the jet was enough to suppress the isoplanatic angles.

## DAY BY DAY SYNOPTIC WEATHER SUMMARY

ANDERSON MESA, FLAGSTAFF, ARIZONA (1989)

Dates: 12 Nov, 2015 hrs - 19 Nov, 0000 hrs (local time)  
13 Nov. 0315 hrs - 19 Nov, 0700 hrs (UTC)

13 November 1989: The synoptic weather pattern over the site is characterized by a surface High; a shortwave propagating through a mid-continental trough (850 mb). At the 700 mb level, the circulation is dictated by the western edge of the mid-continental, longwave trough. From 500 to 200 mb, a Low centered over the Mexico/Arizona border dominates the circulation. By 1200 UTC, this cutoff Low merges with the mid-continental trough (300 mb level).

14 November 1989: The synoptic weather pattern over the site continues with a surface High; a weak shortwave propagating through the mid-country trough (850-700 mb). At 500 mb, the site is west of the longwave trough traversing mid-country. Aloft (0000 UTC), the site is south of a southward moving mid-country trough. By 1200 UTC, the approaching trough moves a 70 kt jet within 5-8 degrees latitude north and south of the site.

15 November 1989: Synoptic weather conditions include a surface 'trof' passing over the site (0000 UTC) with a cold front just to the north. By 1200 UTC, the cold front is located slightly south of the site. At the 850 mb level, the site transitions from the western edge of the mid-country Low (0000 UTC) to the clockwise circulation of the northern Pacific High spreading over Nevada. The 700-500 mb level is dominated by the northwesterly flow around the mid-country trough. Aloft (300-200 mb), a 70-110 kt west-northwesterly jet passes within 2-4 degrees latitude (on all except the southwestern side) of the site.

16 November 1989: Extensive cloud cover limits the optical data sampling. Limited synoptic wind data is available for this session. From 850 to 300 mb, the site is under the western edge of the mid-country trough; the North Pacific High extends over Nevada (850 mb). At 200 mb, the 70-110 kt jet is within 5-8 degrees latitude northwest and southeast of the site.



17 November 1989: The site borders a southeastward moving surface High centered in Oregon and a surface Low over Mexico. Between 850 and 700 mb, the site is on the edge of the extended northern Pacific High/mid-continental trough. At 500 mb, an eastward moving shortwave puts the site under a weak Low. Aloft, a 70 to 110 kt jet is within 5-8 degrees latitude northwest, west and southeast of the site. Though the jet does not traverse the site, the shear set up by the jet's location is enough to suppress the isoplanatic angles.

18 November 1989: The synoptic weather scenario places the site under a southeastward moving surface High centered in the northwestern states and a surface Low over Mexico. At 850 mb, the site is on the edge of the northern Pacific High extension and a flow through the mid-continental trough. From 700-500 mb, an eastward moving shortwave places the site under a weak Low. This Low organizes between 0000 and 1200 UTC. Aloft, a 70 to 110 kt north-northeasterly jet circumnavigates the site (passing within 5-8 degrees latitude). Though the jet does not initially traverse the site, the 200 mb shear set up by the 0000 UTC pattern is approximately  $4.3\text{E-}5 \text{ s}^{-1}$ . By 1200 UTC, a 115 kt northwesterly jet passes over the site. The 200 mb wind shear consequently increases to about  $8.5\text{E-}5 \text{ s}^{-1}$  (very strong shear!!).

19 November 1989: The synoptic weather pattern includes a High pressure system associated with the northern Pacific High extending over Utah. The presence of this High is observed from the surface to 500 mb. At 500 mb, the site borders the High just described and a Low over Baja, California. Aloft, the site is directly under a 70-90 kt northeasterly jet supporting a Low over Mexico. During the 0000-1200 UTC time interval, this Low appears to develop as a cut-off Low over Mexico/southwestern United States. Wind shear over the site at 200 mb (0000 and 1200 UTC) is about  $2.1\text{E-}5 \text{ s}^{-1}$ ; weaker than 18 November, yet still significant.

## APPENDIX B. NWS 850 AND 200 MB SYNOPTIC WEATHER CHARTS

The following National Weather Service (NWS) 850 and 200 mb isobaric weather charts display the synoptic activity present during the 13-19 November 1989 UTC optical measurement session. A plus symbol identifies the location of the Anderson Mesa (Flagstaff, AZ) data collection site. Surface fronts, as well as surface High and Low pressure systems (circled "H" and "L" labels) have been superimposed onto the 850 mb charts. Dashed lines on the 200 mb charts trace the jet stream activity. Specifically, the dashed lines outline the 70 kt isotaches, as well as the labelled jet streak maxima (generally, 110 kt or greater).





Fig 8. NWS 850 (bottom) and 200 (top) mb Charts: November 13, 0000 UTC





Fig 9. NWS 850 (bottom) and 200 (top) mb Charts: November 13, 1200 UTC





Fig 10. NWS 850 (bottom) and 200 (top) mb Charts: November 14, 0000 UTC





Fig 11. NWS 850 (bottom) and 200 (top) mb Charts: November 14, 1200 UTC





Fig 12. NWS 850 (bottom) and 200 (top) mb Charts: November 15, 0000 UTC





Fig 13. NWS 850 (bottom) and 200 (top) mb Charts: November 15, 1200 UTC







Fig 14. NWS 850 (bottom) and 200 (top) mb Charts: November 16, 0000 UTC







Fig 15. NWS 850 (bottom) and 200 (top) mb Charts: November 17, 0000 UTC



**Fig 16. NWS 850 (bottom) and 200 (top) mb Charts: November 17, 1200 UTC**







Fig 17. NWS 850 (bottom) and 200 (top) mb Charts: November 18, 0000 UTC



**Fig 18. NWS 850 (bottom) and 200 (top) mb Charts: November 18, 1200 UTC**





Fig 19. NWS 850 (bottom) and 200 (top) mb Charts: November 19, 0000 UTC






Fig 20. NWS 850 (bottom) and 200 (top) mb Charts: November 19, 1200 UTC

## APPENDIX C. PROCESSED OPTICAL DATA (1989 November 13-19)

Appendix C displays nightly plots of all processed Transverse Coherence Length and Isoplanatic Angle data acquired between 13-19 November 1989.



ANDERSON MESA, AZ - 1989 NOVEMBER 13  
Transverse Coherence Length



Isoplanatic Angle





Fig 21. Anderson Mesa, Az Optical Data: 1989 November 13

ANDERSON MESA, AZ - 1989 NOVEMBER 14  
Transverse Coherence Length



Isoplanatic Angle





Fig 22. Anderson Mesa, Az Optical Data: 1989 November 14

ANDERSON MESA, AZ - 1989 NOVEMBER 15  
Transverse Coherence Length



Isoplanatic Angle





Fig 23. Anderson Mesa, Az Optical Data: 1989 November 15

ANDERSON MESA, AZ - 1989 NOVEMBER 16  
Transverse Coherence Length



Isoplanatic Angle





Fig 24. Anderson Mesa, Az Optical Data: 1989 November 16

ANDERSON MESA, AZ - 1989 NOVEMBER 17  
Transverse Coherence Length



Isoplanatic Angle





Fig 25. Anderson Mesa, Az Optical Data: 1989 November 17

ANDERSON MESA, AZ - 1989 NOVEMBER 18  
Transverse Coherence Length



Isoplanatic Angle





Fig 26. Anderson Mesa, Az Optical Data: 1989 November 18

ANDERSON MESA, AZ - 1989 NOVEMBER 19  
Transverse Coherence Length



Isoplanatic Angle




Fig 27. Anderson Mesa, Az Optical Data: 1989 November 19


#### APPENDIX D. TRANSVERSE COHERENCE LENGTH STATISTICS

Appendix D presents the transverse coherence length ( $r_0$ ) un-normalized percent frequency distribution for each observing night (bin interval is 10 mm; optical wavelength is 500 nm). Empirical seeing quality histograms are also included. The bin intervals selected for this qualitative interpretation are a product of approximately 50 site surveys spanning 18-40 degrees latitude and 65-156 degrees longitude. The specific empirical seeing quality intervals are:

<u>Empirical Seeing Quality</u>	<u><math>r_0</math> measurement (mm)</u>
Poor	00 - 50
Mediocre	51 - 100
Good	101 - 200
Very Good	201 - 300
Excellent	301 - 500



ANDERSON MESA, AZ - 1989 NOVEMBER 13  
 $r_o$  Percent Frequency Distribution



Empirical Seeing Quality




Fig 28. Anderson Mesa, AZ  $r_o$  Statistics: 1989 Nov 13

ANDERSON MESA, AZ - 1989 NOVEMBER 14  
 $r_o$  Percent Frequency Distribution






Fig 29. Anderson Mesa, Az  $r_o$  Statistics: 1989 Nov 14

ANDERSON MESA, AZ - 1989 NOVEMBER 15  
r. Percent Frequency Distribution



Empirical Seeing Quality





Fig 30. Anderson Mesa, Az r. Statistics: 1989 Nov 15

ANDERSON MESA, AZ - 1989 NOVEMBER 16  
 $r_o$  Percent Frequency Distribution



Empirical Seeing Quality





Fig 31. Anderson Mesa, Az  $r_o$  Statistics: 1989 Nov 16

ANDERSON MESA, AZ - 1989 NOVEMBER 17  
 $r_o$  Percent Frequency Distribution



Empirical Seeing Quality





Fig 32. Anderson Mesa, Az  $r_o$  Statistics: 1989 Nov 17

ANDERSON MESA, AZ - 1989 NOVEMBER 18  
*r<sub>0</sub>* Percent Frequency Distribution



Empirical Seeing Quality





Fig 33. Anderson Mesa, Az *r<sub>0</sub>* Statistics: 1989 Nov 18

ANDERSON MESA, AZ - 1989 NOVEMBER 19  
r. Percent Frequency Distribution



Empirical Seeing Quality




Fig 34. Anderson Mesa, Az r. Statistics: 1989 Nov 19


## APPENDIX E. ISOPLANATIC ANGLE STATISTICS

To assist with the interpretation of the isoplanatic angle ( $\theta_0$ ) measurements, an un-normalized frequency distribution and an empirical seeing quality plot for each sampling session have been provided in Appendix E. The bin-size for this frequency distribution is 1 urad. The empirical seeing quality graphs use the following bin intervals:

<u>Empirical Seeing Quality</u>	<u><math>\theta_0</math> measurement (urad)</u>
Very Poor	0 - 4.0
Poor	4.1 - 8.0
Mediocre	8.1 - 12.0
Good	12.1 - 20.0
Very Good	20.1 - 30.0
Excellent	30.1 - 50.0



ANDERSON MESA, AZ - 1989 NOVEMBER 13  
 60 Percent Frequency Distribution



Empirical Seeing Quality





Fig 35. Anderson Mesa, Az  $\theta_0$  Statistics: 1989 Nov 13

ANDERSON MESA, AZ - 1989 NOVEMBER 14  
60 Percent Frequency Distribution



Empirical Seeing Quality





Fig 36. Anderson Mesa, Az  $\theta_0$  Statistics: 1989 Nov 14

ANDERSON MESA, AZ 1989 NOVEMBER 15  
 $\theta_0$  Percent Frequency Distribution



Empirical Seeing Quality




Fig 37. Anderson Mesa, Az  $\theta_0$  Statistics: 1989 Nov 15





Fig 38. Anderson Mesa, Az  $\theta_0$  Statistics: 1989 Nov 16

ANDERSON MESA, AZ - 1989 NOVEMBER 17  
 $\theta_0$  Percent Frequency Distribution






Fig 39. Anderson Mesa, Az  $\theta_0$  Statistics: 1989 Nov 17

ANDERSON MESA, AZ - 1989 NOVEMBER 18  
60 Percent Frequency Distribution



Empirical Seeing Quality





Fig 40. Anderson Mesa, Az 60 Statistics: 1989 Nov 18

ANDERSON MESA, AZ - 1989 NOVEMBER 19  
 $\theta_0$  Percent Frequency Distribution



Empirical Seeing Quality




Fig 41. Anderson Mesa, AZ  $\theta_0$  Statistics: 1989 Nov 19

## APPENDIX F. SEPTEMBER/NOVEMBER $r_0$ AND $\theta_0$ DISTRIBUTION

Appendix F presents the cumulative September and November 1989 normalized frequency distribution for both  $r_0$  and  $\theta_0$ . The measurements incorporated into these figures represent all the 17-28 September and 13-19 November 1989 processed NPS data taken at Anderson Mesa and the United States Naval Observatory near Flagstaff, Arizona. Specifically, Figure 42 displays the distribution generated from the 2,773 September and 2,103 November  $r_0$  samples. The 30,549  $\theta_0$  samples shown in Figure 43 includes: 16,355 September and 14,194 November individual angles.



CUMULATIVE NORMALIZED  $r_0$  FREQUENCY DISTRIBUTION  
1989 September and November - Anderson Mesa, AZ




Fig 42. Cumulative  $r_0$  Distribution: 1989 Sept and Nov. The peak  $r_0$  bin interval is 80-90 mm.

CUMULATIVE NORMALIZED  $\theta_0$  FREQUENCY DISTRIBUTION  
1989 September and November - Anderson Mesa, AZ




Fig 43. Cumulative  $\theta_0$  Distribution: 1989 Sept and Nov. The peak  $\theta_0$  bin interval is 5-6 urad.

## LIST OF REFERENCES

Stevens, K.B., 1985: Remote Measurement of the Atmospheric Isoplanatic Angle and Determination of Refractive Turbulence Profiles by Direct Inversion of the Scintillation Amplitude Covariance Function with Tikhonov Regularization, Ph.D. Dissertation, Naval Postgraduate School, Monterey, CA, 170 pp.

Vaucher, G. Tirrell, 1989: Correlation of Atmospheric Optical Turbulence and Meteorological Measurements, M.S., Naval Postgraduate School, Monterey, CA, 145 pp.

Vaucher, G. Tirrell, Vaucher, Christopher A., Walters, Donald L., 1990: Optical Turbulence and Rawinsonde Measurements for 17-28 September 1989 at Anderson Mesa/United States Naval Observatory, Flagstaff, Arizona, Project Report No. NPS-61-90-005PR, Naval Postgraduate School, Monterey, CA, 105 pp.

Walters, D.L., Favier, D.L., and Hines, J.R., 1979: Vertical Path Atmospheric MTF Measurements. J. Opt. Soc. Am., v. 69, 829.

WeatherMeasure WEATHERtronics, 1987: 1987-1988 Catalog, Geophysical Instruments and Systems, Division of QUALIMETRICS, INC., Sacramento, CA, 323 pp.

# INITIAL DISTRIBUTION LIST

	No. Copies
1. Defense Technical Information Center Cameron Station Alexandria, VA 22304-6145	2
2. Library, Code 52 Naval Postgraduate School Monterey, CA 93943-5002	2
3. Gail Tirrell Vaucher, Code PH Department of Physics Naval Postgraduate School Monterey, CA 93943-5000	6
4. Christopher Vaucher, Code PH USRA Visiting Scientist Department of Physics Naval Postgraduate School Monterey, CA 93943-5000	4
5. Professor Donald L. Walters, Code PH/WE Naval Postgraduate School Monterey, CA 93943-5000	4
6. Dr. Ken Johnston, Code 4130 Naval Research Laboratory Washington, D.C. 20375	1
7. Lowell Observatory Attn: Dr. Bob Millis, Dr. Nat White Mars Hill Rd, 1400 West Flagstaff, AZ 86001	2
8. United States Naval Observatory Attn: Dr. Harold Ables P.O. Box 1149 Flagstaff, AZ 86002	1
9. Dr. Gert Westerhout, Scientific Director United States Naval Observatory 34 Massachusetts Ave, NW Washington, D.C. 20392	1

- |     |  |   |
|-----|--|---|
| 10. | Dr. Frank Kerr<br>Universities Space Research Association<br>Mail Stop 610.3<br>Goddard Space Flight Center<br>Greenbelt, Maryland 20771 | 1 |
| 11. | Dr. Robert Fugate<br>WL/ARCA<br>Kirtland AFB, New Mexico 87117-6002  | 1 |
| 12  | Capt. Ann Slavin<br>WL/ARCA<br>Kirtland AFB, New Mexico 87117-6002   | 1 |
| 13. | Dean of Research, Code 08<br>Naval Postgraduate School<br>Monterey, CA 93943-5000  | 1 |
| 14. | Department of Physics, Code PH<br>Naval Postgraduate School<br>Monterey, CA 93943-5000   | 1 |